



A Question Mark on Emerging Zinc-Related Nutritional Deficiencies in Pakistani Population

Author(s): Waqas Asghar, Wahab Nazir and Nauman Khalid

Source: *Asia Pacific Journal of Public Health*, July 2018, Vol. 30, No. 5 (July 2018), pp. 500-502

Published by: Sage Publications, Inc.

Stable URL: <https://www.jstor.org/stable/10.2307/27009129>

REFERENCES

Linked references are available on JSTOR for this article:

https://www.jstor.org/stable/10.2307/27009129?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



Sage Publications, Inc. is collaborating with JSTOR to digitize, preserve and extend access to *Asia Pacific Journal of Public Health*

JSTOR



A Question Mark on Emerging Zinc-Related Nutritional Deficiencies in Pakistani Population

**Waqas Asghar, BSc, MSc¹, Wahab Nazir, BSc, MSc¹,
and Nauman Khalid, MSc, PhD¹** 

To the Editor:

Micronutrient deficiencies, often referred to as “hidden hunger,” are a critically significant, though often an imperceptible form of malnutrition¹ in various population segments of developing countries such as Pakistan. Zinc, a trace element, is a key component for the effective functioning of nearly 300 metalloenzymes, as well as having roles in metabolism mechanisms of DNA and RNA, immune response mechanisms, signal transduction, cell division, and gene expression.^{2,3} Zinc deficiency in pregnant women, if untreated, can have significant negative pregnancy outcomes and linear growth problems for children, such as premature delivery, childhood diarrhea–associated morbidity, acute lower respiratory tract infections, stunting, wasting, and underweight children.^{4,5}

Serum zinc concentration studies are the key indicators and mainstay of determination of zinc deficiency levels in Pakistan. Regional and nationwide surveys and biochemical analyses in Pakistan indicate that the subclinical deficiency of zinc among pregnant as well as nonpregnant women and children is widely prevalent, with the most recent National Nutrition Survey of 2011 reporting zinc deficiency in 47.6% of pregnant women, 41.3% nonpregnant women, and 39.2% children under the age of 5 years (Figure 1).⁶ Moreover, the survey also indicated that the prevalence of zinc deficiency did not experience a major shift, with 41.9% reported in National Nutrition Survey of 2009 and 41.3% in its updated volume of 2011; however, there was a 2.1% increase in zinc deficiency among children under 5 years of age over the same period.⁶

The findings regarding zinc deficiency in Pakistan have also been substantiated by international research on this public health challenge. Akhtar⁴ reported that a few studies conducted on zinc deficiency in Pakistan indicated 54.2% school-going children and 37.1% children of preschool age to be deficient. Moreover, nearly half of children under 5 years of age are stunted, and about 1 in every 3 is underweight in Pakistan.¹ However, currently, the key limitation in determining the precise extent and prevalence of zinc deficiency in Pakistan appears to be the insufficient number of studies being conducted for the evaluation and monitoring of the situation over the past 4 to 6 years. This trend, in turn, is also hampering the mitigation efforts and formulation of effective implementation strategies for the immediate and long-term future scenarios.

¹University of Management and Technology, Lahore, Pakistan

Corresponding Author:

Nauman Khalid, School of Food and Agricultural Sciences, University of Management and Technology, Lahore 54000, Pakistan.

Email: nauman.khalid@umt.edu.pk




Figure 1. Zinc availability in national food supplies and the prevalence of stunting. The prevalence of zinc deficiency in Pakistani children less than 5 years of age.^{1,6} The red line indicates significant zinc deficiency in almost all provinces of Pakistan. KP stands for Khyber Pukhtunkhwa provinces, AJK stands for Azad Jumma and Kashmir, and GB stands for Gilgit-Baltistan.

It has been widely acknowledged over the decades that micronutrient deficiencies are highly prevalent across Pakistan. But as Wieser et al¹ reported, apart from policies for the eradication of iodine deficiency, the improvement in deficiency levels for other micronutrients has been largely insignificant. Likewise, the guidelines set forth by the Pakistan Standards and Quality Control Act for the mass fortification of essential foodstuff such as wheat flour, edible oil, and ghee, as well as iodized salt, have been, except for iodized salt, mostly unsuccessful.¹ Therefore, effective execution of policies such as complementary feeding, diet diversification, mass fortification, and multiple micronutrient supplementation, aimed at both pregnant and breastfeeding mothers, as well as young school-age children, should be the way forward.

ORCID iD

Nauman Khalid  <https://orcid.org/0000-0002-8045-199X>

References

1. Wieser S, Brunner B, Tzogiou C, et al. Societal costs of micronutrient deficiencies in 6- to 59-month-old children in Pakistan. *Food Nutr Bull.* 2017;38:485-500.
2. Kumera G, Awoke T, Melese T, et al. Prevalence of zinc deficiency and its association with dietary, serum albumin and intestinal parasitic infection among pregnant women attending antenatal care at the University of Gondar Hospital, Gondar, Northwest Ethiopia. *BMC Nutr.* 2015;1:31. doi:10.1186/s40795-015-0026-6.
3. Khalid N, Ahmed A, Bhatti MS, Randhawa MA, Ahmad A, Rafaqat R. A question mark on zinc deficiency in 185 million people in Pakistan—possible way out. *Crit Rev Food Sci Nutr.* 2014;54:1222-1240.
4. Akhtar S. Zinc status in South Asian populations—an update. *J Health Popul Nutr.* 2013;31:139-149.
5. Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One.* 2012;7:e50568.
6. Planning Commission (Planning and Development Division). *National Nutrition Survey.* Islamabad, Pakistan: Government of Pakistan; 2011.