



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

Solving directly third-order ODEs using operational matrices of Bernstein polynomials method with applications to fluid flow equations

Sana'a Nazmi Khataybeh^{a,*}, Ishak Hashim^a, Mohammed Alshbool^b^a School of Mathematical Sciences, Faculty of Science & Technology, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia^b Faculty of Art & Sciences, Abu Dhabi University, Abu Dhabi, United Arab Emirates

ARTICLE INFO

Article history:

Received 1 September 2017

Accepted 1 May 2018

Available online 5 May 2018

Keywords:

ODEs

Bernstein polynomials

Approximation

Fluid flow

ABSTRACT

In this paper, we adapt for the first time the operational matrices of Bernstein polynomials method for solving directly a class of third-order ordinary differential equations (ODEs). This method gives a numerical solution by converting the equation into a system of algebraic equations which is solved directly. Applications of the present method to the famous Blasius equation describing a boundary layer flow over a flat plate and third-order ODE for thin film flow are presented. Some numerical examples are also given to show the applicability of the method.

© 2018 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The mathematical formulations of physical phenomena and engineering often lead to ordinary differential equations (ODEs). Finding an approximate analytical solution to an equation has attracted the attention of many researchers (see for example Zhang et al., 2016; Singh et al., 2017; Kumar et al., 2017). There are currently many methods that can provide approximate solutions, and most of these methods first convert the ODEs to a system of lower degree, usually the first-order.

In recent years, several authors have introduced direct methods for solving higher-order ODEs. Majid et al. (2009) used Jacobi iteration and direct methods with a variable step size for solving second-order ODEs. Awoyemi and Idowu (2005) proposed a class of hybrid collocation direct method to solve third-order ODEs. A part from that, power series collocation and interpolation was implemented to derive a 3-step block method for solving ODEs of the third-order by Olabode and Yusuph (2009). Waeleh et al. (2011) developed a block method based on numerical integration and interpolation in order to solve higher-order ODEs where they

presented approximation solutions of the fourth- and fifth-orders. Olabode and Alabi (2013) introduced a linear multistep method using interpolation and collocation of power series approximation solution to solve the fourth-order ODEs.

There have been quite a number of papers on applying Bernstein polynomials for solving differential equations. Bernstein polynomials were introduced by Sergei Bernstein in 1912 (see Lorentz, 2012). Yousefi and Behroozifar (2010) employed operational matrices of Bernstein polynomials for solving differential equations. Pandey and Kumar (2012) used Bernstein operational matrices to solve Emden type equations. Alshbool et al. (2015) introduced approximation solutions for singular differential equations using Bernstein polynomials. A numerical solution for the variable order linear cable equation using Bernstein polynomials was presented by Chen et al. (2014). Bellucci (2014) introduced the orthonormal Bernstein polynomials which can be used in a generalized Fourier series to approximate surfaces and curves. Mirkov and Rašuo (2013), Mirkov et al., 2012 used Bernstein polynomials to solve elliptic boundary value problems, in particular, the Poisson and Helmholtz equations on a square domain. Further capabilities of operational matrices of Bernstein polynomials were demonstrated very recently by Asgari and Ezzati (2017) who solved two-dimensional integral equations of fractional order by operational matrix of two-dimensional Bernstein polynomials and Alshbool et al. (2017) who solved fractional-order differential equations. Very recently, Loh et al. (2017) introduced a new operational matrix based on Genocchi polynomials for solving fractional integro-differential equations.

* Corresponding author.

E-mail address: sanaakh153@gmail.com (S.N. Khataybeh).

Peer review under responsibility of King Saud University



Production and hosting by Elsevier

In this paper, we introduce for the first time an approximate method based on Bernstein polynomials for solving directly third-order ODEs of the form

$$y''' = f(x, y, y', y'') \quad \text{subject to} \quad y(0) = a_0, y'(0) = b_0, y'' = c_0, \quad (1)$$

where a_0, b_0, c_0 are given constants and $x \in [0, 1]$. While all others methods as introduced in Zhang et al. (2016), Singh et al. (2017), Kumar et al. (2017), Majid et al. (2009), Awoyemi and Idowu (2005)) solved Eq. (1) by reducing it to systems of lower degree of ODEs, the present method converts Eq. (1) to a system of algebraic equations which can be solved easily. The capability of the method shall be tested on a linear and nonlinear third-order ODEs.

2. Description of method

The Bernstein polynomials of degree n are defined as

$$B_{v,n}(x) = \binom{n}{v} x^v (1-x)^{n-v}, \quad (2)$$

where

$$\binom{n}{v} = \frac{n!}{v!(n-v)!}, \quad v = 0, \dots, n. \quad (3)$$

These polynomials form a complete basis for the vector space Π_n of polynomials of degree at most n . For convenience, $B_{v,n}(x) = 0$ if $v < 0, n < v$. These polynomials have many properties which make them important and useful, like continuity and unity partition (see Farouki, 2012). As a result, any polynomial of degree n can be approximated in terms of linear combination of $B_{v,n}(x) = 0, (v = 0, \dots, n)$ as given below,

$$y(x) = \sum_{v=0}^n C_v B_{v,n} = C^T \Phi(x), \quad (4)$$

where $C^T = [C_0, C_1, \dots, C_n]$ and $\Phi(x) = [B_{0,n}, B_{1,n}, \dots, B_{n,n}]^T$.

Also, we can make the decomposition of the vector $\Phi(x)$ as a product of a square matrix of size $(n+1) \times (n+1)$ and a vector of size $(n+1) \times 1$, i.e. $\Phi(x) = AX$, where

$$A = \begin{bmatrix} a_0 & a_1 & a_2 & \dots & a_n \\ b_0 & b_1 & b_2 & \dots & b_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ k_0 & k_1 & k_2 & \dots & k_n \end{bmatrix}, \quad X = \begin{bmatrix} 1 \\ x \\ x^2 \\ \vdots \\ x^n \end{bmatrix}. \quad (5)$$

Now, the derivative of the vector $\Phi(x)$ denoted by D' is

$$\frac{d\Phi(x)}{dx} = D'\Phi(x), \quad (6)$$

where D' is the $(n+1) \times (n+1)$ operational matrix of the derivative that given as $D' = A\sigma A^{-1}$, where

$$A = \begin{bmatrix} a_0 & a_1 & a_2 & \dots & a_n \\ b_0 & b_1 & b_2 & \dots & b_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ k_0 & k_1 & k_2 & \dots & k_n \end{bmatrix}, \quad \sigma = \begin{bmatrix} 0 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & 0 \end{bmatrix}. \quad (7)$$

for more illustration, (see Yousefi and Behroozifar, 2010). Eq. (6) can be generalised as follows

$$\frac{d^2\Phi(x)}{dx^2} = (D')^2\Phi(x), \dots, \frac{d^n\Phi(x)}{dx^n} = (D')^n\Phi(x). \quad (8)$$

To use this operational matrix for solving an equation, we will approximate $y(x)$ by Bernstein polynomials as $y(x) = C^T\Phi(x)$ and we have

$$y'(x) = C^T D' \Phi(x), \quad y''(x) = C^T (D')^2 \Phi(x), \quad y'''(x) = C^T (D')^3 \Phi(x). \quad (9)$$

Then Eqn. (1) can be written in the form

$$C^T (D')^3 \Phi(x) = f(x, C^T \Phi(x), C^T D' \Phi(x), C^T (D')^2 \Phi(x)), \quad (10)$$

$$C^T \Phi(x_0) = a_0, \quad C^T D' \Phi(x_0) = b_0, \quad C^T (D')^2 \Phi(x_0) = c_0. \quad (11)$$

We substitute the collocation nodes $0 \leq x_0 < x_1 < \dots < x_m \leq 1$ in (10) to get a system of algebraic equations which will be solved by a computer algebra system like Maple. We use the Chebyshev roots as the collocation nodes

$$x_i = \frac{1}{2} + \frac{1}{2 \cos((2i+1)\frac{\pi}{2n})}, \quad i = 0, 1, \dots, n-2. \quad (12)$$

A convergent analysis of the method was given by Yousefi et al. (2011) in their lemmas 3 and 4. For more details the readers is referred to Rivlin (2003).

3. Numerical tests

In this section, we present examples of third-order ODEs in order to illustrate the performance and effectiveness of our method. We apply the method with the number of Bernstein terms, $n = 14$. All the algebraic manipulations were done in Maple 8 with the Digits set to 20.

3.1. Example 1

First we consider the following linear problem

$$y''' - y'' + y' - y = 0, \quad y(0) = 1, y'(0) = 0, y''(0) = -1, \quad 0 \leq x \leq 1,$$

with the exact solution $y(x) = \cos x$. Applying the method described above, the following 14-term approximate solution is obtained:

$$\begin{aligned} y(x) \approx & -9.64735391 \times 10^{-12} x^{14} - 9.3279316 \times 10^{-12} x^{13} \\ & + 2.111956063 \times 10^{-9} x^{12} - 4.022855 \times 10^{-11} x^{11} \\ & - 2.755270872610^{-7} x^{10} - 3.7973510^{-11} x^9 \\ & + 0.248016101554 \times 10^{-4} x^8 - 1.00854 \times 10^{-11} x^7 \\ & - 0.13888888856464 \times 10^{-2} x^6 - 7.443 \times 10^{-13} x^5 \\ & + 0.4166666666678504 \times 10^{-1} x^4 - 1.264 \times 10^{-14} x^3 \\ & - 0.50000000000000001 x^2 \\ & + 0.999999999999999999. \end{aligned}$$

The absolute errors shown in Table 1 suggest that the present method is very accurate.

Table 1
Absolute errors for Example 1.

x	Exact solution	Bernstein solution	Error
0.1	0.99500416527802576610	0.99500416527802576026	5.84×10^{-18}
0.2	0.98006657784124163112	0.98006657784124160265	2.85×10^{-17}
0.3	0.95533648912560601964	0.95533648912560595103	6.86×10^{-17}
0.4	0.92106099400288508280	0.92106099400288495944	1.23×10^{-16}
0.5	0.87758256189037271612	0.87758256189037253350	1.83×10^{-16}
0.6	0.82533561490967829724	0.82533561490967807361	2.24×10^{-16}
0.7	0.76484218728448842626	0.76484218728448822968	1.97×10^{-16}
0.8	0.69670670934716542092	0.69670670934716545101	3.01×10^{-17}
0.9	0.62160996827066445648	0.62160996827066530979	8.53×10^{-16}
1.0	0.54030230586813971740	0.54030230586814316648	3.45×10^{-15}




Fig. 1. Approximation solution for example 1 using Bernstein polynomials with exact solution.

3.2. Example 2

Next consider the following problem [Fig. 1](#)

$$y''' + 5y'' + 7y' + 3y = 0, \quad y(0) = 1, \quad y'(0) = 0, \\ y''(0) = -1, \quad 0 \leq x \leq 1.$$

The exact solution is $y(x) = e^{-x} + xe^{-x}$. The operational matrices of Bernstein polynomials method yields

$$y(x) \approx -8.023524662 \times 10^{-11}x^{14} + 1.683945475 \times 10^{-9}x^{13} \\ - 2.2420566212 \times 10^{-8}x^{12} + 2.4968885757 \times 10^{-7}x^{11} \\ - 0.247924992485 \times 10^{-5}x^{10} + 0.220451309070 \times 10^{-4}x^9 \\ - 0.1736106853770 \times 10^{-3}x^8 + 0.11904760061469 \\ \times 10^{-2}x^7 - 0.69444443862206 \times 10^{-2}x^6 \\ + 0.33333333201911 \times 10^{-1}x^5 \\ - 0.12499999999796276x^4 + 0.33333333333314344x^3 \\ - 0.500000000000000001x^2 \\ + 0.999999999999999999.$$

[Table 2](#) clearly shows the accuracy of the present method.

Table 2
Absolute errors for Example 2.

x	Exact solution	Bernstein solution	Error Present method	Error Ref. (Mohammed and Adeniyi, 2014)
0.1	0.9953211598395553048	0.99532115983955545628	7.42×10^{-17}	1.00×10^{-10}
0.2	0.98247690369357823040	0.98247690369357792919	3.01×10^{-16}	3.00×10^{-10}
0.3	0.96306368688623322589	0.96306368688623261523	6.11×10^{-16}	7.00×10^{-10}
0.4	0.93844806444989502104	0.93844806444989407304	9.48×10^{-16}	7.00×10^{-10}
0.5	0.90979598956895013540	0.90979598956894887800	1.26×10^{-15}	6.00×10^{-10}
0.6	0.87809861775044229221	0.87809861775044086917	1.42×10^{-15}	2.00×10^{-10}
0.7	0.84419501644539617499	0.84419501644539506014	1.11×10^{-15}	9.00×10^{-10}
0.8	0.80879213541099886457	0.80879213541099944217	5.78×10^{-16}	2.80×10^{-9}
0.9	0.77248235350713831257	0.77248235350714424660	5.93×10^{-15}	5.40×10^{-9}
1.0	0.73575888234288464320	0.73575888234290481537	2.02×10^{-14}	3.50×10^{-9}

3.3. Example 3

$$y''' = 3 \sin x, \quad y(0) = 1, \quad y'(0) = 0, \quad y''(0) = -2, \quad 0 \leq x \leq 1.$$

The exact solution of this problem is $y(x) = 3 \cos x + (x^2/2) - 2$ and the approximate solution of the present method is

$$y(x) \approx -2.374911736 \times 10^{-11}x^{14} - 6.86371074 \times 10^{-11}x^{13} \\ + 6.478659491 \times 10^{-9}x^{12} - 4.1795110 \times 10^{-10}x^{11} \\ - 8.2617332888 \times 10^{-7}x^{10} - 5.023725 \times 10^{-10}x^9 \\ + 0.744050936033 \times 10^{-4}x^8 - 1.581103 \times 10^{-10}x^7 \\ - 0.41666666125723 \times 10^{-2}x^6 - 1.30540 \times 10^{-11}x^5 \\ + 0.12500000000215427x^4 - 2.3127 \times 10^{-13}x^3 \\ - 1.000000000000000001x^2 \\ + 0.999999999999999999.$$

Again the errors depicted in [Table 3](#) show the accuracy of the present method. [Figs. 2 and 3](#).

4. Applications to equations of fluid flow

4.1. Boundary layer flow

Now consider the nonlinear boundary layer equation

$$2y''' + yy'' = 0, \quad y(0) = 0, \quad y'(0) = 0, \quad y''(0) = 1. \quad (13)$$

This equation is famously known as the Blasius equation. The aim of solving Blasius equation to get the value $y''(0)$ to evaluate the shear stress at the plate. Blasius equation has been solved using different methods like series expansions, Runge Kutta, differential transformation and others. The interested readers can also see [Lien-Tsai and Cha'o-Kuang \(1998\)](#) and [Schlichting et al. \(1955\)](#).

Table 3
Absolute errors for Example 3.

x	Exact solution	Bernstein solution	Error
0.1	0.9900124958340772983	0.9900124958340771930	1.05×10^{-16}
0.2	0.9601997335237248934	0.96019973352372439108	5.02×10^{-16}
0.3	0.9110094673768180589	0.91100946737681686283	1.20×10^{-15}
0.4	0.8431829820086552484	0.93844806444989407304	2.18×10^{-15}
0.5	0.7577476856711181484	0.84318298200865306768	3.43×10^{-15}
0.6	0.6560068447290348917	0.65600684472902999918	4.89×10^{-15}
0.7	0.5395265618534652788	0.53952656185345885671	6.42×10^{-15}
0.8	0.4101201280414962628	0.41012012804148848222	7.78×10^{-15}
0.9	0.2698299048119933694	0.26982990481198474225	8.63×10^{-15}
1.0	0.1209069176044191522	0.12090691760441048523	8.67×10^{-15}




Fig. 2. Approximation solution for example 2 using Bernstein polynomials with exact solution.




Fig. 3. Approximation solution for example 3 using Bernstein polynomials with exact solution.

$$y(x) \approx 1.12610441 \times 10^{-7}x^{14} - 7.7621579 \times 10^{-7}x^{13} \\ + 0.27970246 \times 10^{-5}x^{12} - 0.7042654 \times 10^{-5}x^{11} \\ + 0.8050211 \times 10^{-5}x^{10} - 0.7633041 \times 10^{-5}x^9 \\ + 0.73343596 \times 10^{-4}x^8 - 0.2480836 \times 10^{-5}x^7 + 8.55631 \\ \times 10^{-7}x^6 - 0.4166874207 \times 10^{-2}x^5 + 3.43594 \times 10^{-8}x^4 \\ - 3.69550 \times 10^{-9}x^3 + 0.5x^2.$$

Comparison with the results from a hybrid block method of Adesanya et al. (2013) shows a good agreement (see Table 4).

4.2. Thin film flow

The motion of the contact line for a thin oil drop spreading on a horizontal surface can be modelled by

Table 4

Comparison with hybrid block method of Adesanya et al. (2013).

x	Ref. (Adesanya et al., 2013)	Present method
0.1	0.00499997916611	0.0049999583341723
0.2	0.0199986666859	0.0199986668419935
0.3	0.044998481293978	0.0449898794745896
0.4	0.079991467388617	0.0799573779857994
0.5	0.124967454367055	0.1248700575229549
0.6	0.179902837409194	0.179677141245484
0.7	0.244755067600357	0.244303616982151
0.8	0.319454500640289	0.318646009310246
0.9	0.403894871267148	0.402568620552525
1.0	0.49792248311043	0.495900382783151

$$y''' = y^{-2}. \quad (14)$$

This equation describes the dynamic balance between surface tension and viscous forces in the absence of gravity in a thin fluid layer. Duffy and Wilson (1997) introduced a parametric representation for the exact solution of Eq. (14) as

$$x = 2^{\frac{1}{3}}\pi \frac{A_i(s)B_i(s_0) - A_i(s_0)B_i(s)}{[aA_i(s_0) + bB_i(s_0)][aA_i(s) + bB_i(s)]}, y = \frac{1}{[aA_i(s) + bB_i(s)]^2},$$

where $A_i(\cdot), B_i(\cdot)$ are the Airy functions, and a, b and s_0 are constants determined by the initial conditions. This equation was also solved by Mechee et al. (2013) and Momoniat and Mahomed (2010). For the initial conditions $y(0) = y'(0) = y''(0) = 1$, Momoniat and Mahomed (2010) obtained

$$a = 0.676482, \quad b = 1.60629, \quad s_0 = -0.39685. \quad (15)$$

We apply our method on this problem with $n = 14$ and Chebyshev roots as a collocation nodes. Applying the present method gives

$$y(x) \approx -0.45984700218196 \times 10^{-5}x^{14} + 0.48584303974767 \\ \times 10^{-4}x^{13} - 0.24134233269265 \times 10^{-3}x^{12} \\ + 0.74592504943532 \times 10^{-3}x^{11} - 0.15918744859666 \\ \times 10^{-2}x^{10} + 0.24661462173050 \times 10^{-2}x^9 \\ - 0.30219812091711 \times 10^{-2}x^8 + 0.43032528580877 \\ \times 10^{-2}x^7 - 0.110923450695375 \times 10^{-1}x^6 \\ + 0.333291876770568 \times 10^{-1}x^5 - 0.833326916924882 \\ \times 10^{-1}x^4 + 0.16666660079020539x^3 \\ + 0.4999999999999999x^2 + x + 1.$$

Comparison in Table 5 shows that our present method yields numerical results which are correct to five decimal places.

Table 5

Solutions of the present method as compared with the exact solution and Runge-Kutta type solution.

x	Exact solution	Ref. (Mechee et al., 2013)	Present method
0.0	1.000000000	1.0000000000	1.0000000000
0.2	1.221211030	1.2212100045	1.2212100043
0.4	1.488834893	1.4888347799	1.4888347792
0.6	1.807361404	1.8073613977	1.8073613962
0.8	2.179819234	2.1798192339	2.1798192314
1	2.608275822	2.6082748676	2.6082748636

5. Conclusion

In this paper, we have modified for the first time the classical operational matrices of Bernstein polynomials method to solve directly a class of third-order ODEs. The method presented in this work avoids the need to transform the ODEs to a system of lower (or first)-order ODEs. The approximate solutions obtained are in the form of series whose terms can be easily computed. The procedure of the method can be programmed symbolically softwares like Maple or Mathematica. The method has also been shown to perform reasonably well for all the test problems.

References

- Adesanya, A.O., Udoh, D.M., Ajileye, A., 2013. A new hybrid block method for the solution of general third order initial value problems of ordinary differential equations. *Int. J. Pure Appl. Math.* 86, 365–375.
- Alshbool, M., Bataineh, A., Hashim, I., Isik, O.R., 2015. Approximate solutions of singular differential equations with estimation error by using Bernstein polynomials. *Int. J. Pure Appl. Math.* 100, 109–125.
- Alshbool, M.H.T., Bataineh, A.S., Hashim, I., Isik, O.R., 2017. Solution of fractional-order differential equations based on the operational matrices of new fractional Bernstein functions. *J. King Saud. Univ. Sci.* 29, 1–18.
- Asgari, M., Ezzati, R., 2017. Using operational matrix of two-dimensional Bernstein polynomials for solving two-dimensional integral equations of fractional order. *Appl. Math. Comput.* 307, 290–298.
- Awoyemi, D., Idowu, O., 2005. A class of hybrid collocation methods for third-order ordinary differential equations. *Int. J. Comput. Math.* 82, 1287–1293.
- Bellucci, M.A., 2014. On the explicit representation of orthonormal Bernstein polynomials. *arXiv preprint arXiv:1404.2293*. Bibliographic Code: arXiv:1404.2293 [math.CA], Cornell University Library. 13 pages.
- Chen, Y., Liu, L., Li, B., Sun, Y., 2014. Numerical solution for the variable order linear cable equation with Bernstein polynomials. *Appl. Math. Comput.* 238, 329–341.
- Duffy, B., Wilson, S., 1997. A third-order differential equation arising in thin-film flows and relevant to Tanner's law. *Appl. Math. Lett.* 10, 63–68.
- Farouki, R.T., 2012. The Bernstein polynomial basis: a centennial retrospective. *Comput. Aided Geom. Des.* 29, 379–419.
- Kumar, S., Kumar, A., Odibat, Z.M., 2017. A nonlinear fractional model to describe the population dynamics of two interacting species. *Math. Methods Appl. Sci.* 40, 4134–4148.
- Lien-Tsai, Y., Cha'o-Kuang, C., 1998. The solution of the Blasius equation by the differential transformation method. *Math. Comput. Model.* 28, 101–111.
- Loh, J.R., Phang, C., Isah, A., 2017. New operational matrix via Genocchi polynomials for solving Fredholm-Volterra fractional integro-differential equations. *Adv. Math. Phys.* Article ID 3821870.
- Lorentz, G.G., 2012. Bernstein polynomials. *Amer. Math. Soc.*, New York.
- Majid, Z.A., Azmi, N.A., Suleiman, M., 2009. Solving second order ordinary differential equations using two point four step direct implicit block method. *Eur. J. Sci. Res.* 31, 29–36.
- Mechee, M., Senu, N., Ismail, F., Nikouravan, B., Siri, Z., 2013. A three-stage fifth-order Runge-Kutta method for directly solving special third-order differential equation with application to thin film flow problem. *Math. Probl. Eng.* 2013, Article ID 795397.
- Mirkov, N., Rašuo, B., 2013. Bernstein polynomial collocation method for elliptic boundary value problems. *PAMM* 13, 421–422.
- Mirkov, N., Rasuo, B., 2012. A Bernstein polynomial collocation method for the solution of elliptic boundary value problems. *arXiv preprint arXiv:1211.3567*. Bibliographic Code: 2012arXiv1211.3567M, Cornell University Library. 21 pages.
- Mohammed, U., Adeniyi, R., 2014. A three step implicit hybrid linear multistep method for the solution of third order ordinary differential equations. *Gen. Math. Notes.* 25, 62–74.
- Momoniat, E., Mahomed, F., 2010. Symmetry reduction and numerical solution of a third-order ode from thin film flow. *Math. Comput. Appl.* 15, 709–719.
- Olabode, B.T., Alabi, T., 2013. Direct block predictor-corrector method for the solution of general fourth order odes. *J. Math. Res.* 5, 26–33.
- Olabode, B., Yusuph, Y., 2009. A new block method for special third order ordinary differential equations. *J. Math. Stat.* 5, 167–170.
- Pandey, R.K., Kumar, N., 2012. Solution of Lane-Emden type equations using Bernstein operational matrix of differentiation. *New Astron.* 17, 303–308.
- Rivlin, T.J., 2003. *An Introduction to the Approximation of Functions*. Courier Corporation, New York.
- Schlichting, H., Gersten, K., Krause, E., Oertel, H., 1955. *Boundary-Layer Theory*, volume 7. Springer, New York.
- Singh, J., Kumar, D., Swroop, R., Kumar, S., 2017. An efficient computational approach for time-fractional Rosenau-Hyman equation. *Neural Comput. Appl.*, 1–8 <https://doi.org/10.1007/s00521-017-2909>.
- Waele, N., Abdul Majid, Z., Ismail, F., Suleiman, M., 2011. Numerical solution of higher order ordinary differential equations by direct block code. *J. Math. Stat.* 8, 77–81.
- Yousefi, S., Behroozifar, M., 2010. Operational matrices of Bernstein polynomials and their applications. *Int. J. Syst. Sci.* 41, 709–716.
- Yousefi, S., Behroozifar, M., Dehghan, M., 2011. The operational matrices of Bernstein polynomials for solving the parabolic equation subject to specification of the mass. *J. Comput. Appl. Math.* 235, 5272–5283.
- Zhang, Y., Kumar, A., Kumar, S., Baleanu, D., Yang, X.J., 2016. Residual power series method for time-fractional Schrödinger equations. *J. Nonlinear Sci. Appl.* 9, 5821–5829.