



## On the Geometry of Weak Fuzzy Complex Numbers and Applications to the Classification of Some A-Curves

Abdallah Shihadeh<sup>1</sup>, Wael Mahmoud M. Salameh<sup>2,\*</sup>, Malik Bataineh<sup>3</sup>, Hassan Al-Tarawneh<sup>4</sup>, Ayman Alahmade<sup>5</sup>, Abdallah Al-Husban<sup>6</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, The Hashemite University, Zarqa 13133, PO box 330127, Jordan

<sup>2</sup>Faculty of Information Technology, Abu Dhabi University, Abu Dhabi, UAE

<sup>3</sup>Department of Mathematics and Statistics, Jordan University of Science and Technology, Irbid, Jordan,

<sup>4</sup>Department of Data sciences and Artificial Intelligence, Al-Ahliyya Amman University, Amman, Jordan

<sup>5</sup>Department of Mathematics, College of Science and Art, AlUla branch, Taibah University, Medina, Saudi Arabia

<sup>6</sup>Department of Mathematics, Faculty of Science and Technology, Irbid National University, P.O. Box: 2600 Irbid, Jordan

Emails: [abdallaha\\_ka@hu.edu.jo](mailto:abdallaha_ka@hu.edu.jo); [wael.salameh@adu.ac.ae](mailto:wael.salameh@adu.ac.ae); [msbataineh@just.edu.jo](mailto:msbataineh@just.edu.jo); [H.Altarawneh@Ammanu.edu.jo](mailto:H.Altarawneh@Ammanu.edu.jo); [aaahmdi@taibahu.edu.sa](mailto:aaahmdi@taibahu.edu.sa); [dralhoshban@inu.edu.jo](mailto:dralhoshban@inu.edu.jo)

### Abstract

The concept of A-curves is considered as a novel application of real field extensions in solving some algebraic vectorial equations defined by Euclidean norms. In this paper, we present a novel insight through the classification of A-curves by illustrating many new semi-module isomorphisms between the direct product of weak fuzzy complex numbers with itself and the direct product of classical Euclidean vector spaces multiplied by itself. These isomorphisms will give us a full classification of A-curves that are related to weak fuzzy complex ring. Also, we provide many examples to explain the contribution of our work.

**Keywords:** weak fuzzy complex number; weak fuzzy complex vector space; A-curve; vectorial equation.

### 1. Introduction

Algebraic extensions of real numbers have always been the focus of attention of researchers from everywhere in the world, where in previous research we find concepts dedicated to the construction of extended algebraic groups and rings of real numbers, such as the split-complex numbers, weak fuzzy numbers, neutrosophic numbers, and even n-cyclic numbers [1-2, 8-9].

The weak fuzzy numbers were defined as a new ring expansion of the real numbers [3], defined by the following formula:  $F_J = \{x_0 + x_1 J; x_0, x_1 \in \mathbf{R}, J^2 = t \in ]0, 1[ \}$

Let  $X = x_0 + x_1 J, Y = y_0 + y_1 J \in F_J$ , where  $x_0, x_1, y_0, y_1 \in \mathbf{R}$

Addition operation:  $X + Y = (x_0 + y_0) + (x_1 + y_1)J$ .

Multiplication operation:  $X \cdot Y = (x_0 y_0 + x_1 y_1 t) + (x_0 y_1 + x_1 y_0)J$ .

These numbers have been studied by many researchers, where in [10,12] we find outstanding efforts to find weak fuzzy Pythagorean triples and quadruples, and solutions to the nonlinear Diophantine equations associated with them. In [13], the computer was programmed to deal with these numbers through the Python environment as a direct application in Computer Science.

Also in [4-5], we find an algebraic study of spaces related to weak fuzzy numbers and applications of inner products to them. In [11], the researchers studied some vector equations defined by the Euclidean norms in Euclidean inner product spaces, where these equations resulted in various algebraic surfaces called A-curves.

As an open research question, the researchers wondered about the possibility of classifying these surfaces by developing the tools used in the study of weak fuzzy numbers.

All this prompted us to present a novel insight through the classification of A-curves by illustrating many new semi-module isomorphisms between the direct product of weak fuzzy complex numbers with itself and the direct product of classical Euclidean vector spaces multiplied by itself. These isomorphisms will give us a full classification of A-curves that are related to weak fuzzy complex ring.

## 2. Main discussion

### Definition:

The ring of weak fuzzy complex numbers is defined as follows.

$$W = \{a + bJ; J^2 = t \in ]0,1[; a, b \in R\}$$

Addition on W is defined as follows:

$$(a + bJ) + (c + dJ) = (a + c) + (b + d)J,$$

Multiplication on W is defined as follows:

$$(a + bJ)(c + dJ) = (ac + bdt) + (ad + bc)J; J^2 = t \in ]0,1[.$$

### Definition:

Let  $W^2 = \{(x, y); x, y \in W\}$ , define

$$(+)\ W^2 \times W^2 \rightarrow W^2; (x_0, x_1) + (y_0, y_1) = (x_0 + y_0, x_1 + y_1)$$

$$(\cdot)\ W^2 \times W^2 \rightarrow W^2; (x_0, x_1) \cdot (y_0, y_1) = (x_0 y_0, x_1 y_1)$$

$$(\times)\ W \times W^2 \rightarrow W^2; (a + bJ) \times (x_0, x_1) = ((a + bJ)x_0, (a + bJ)x_1),$$

Where  $x_0, x_1, y_0, y_1 \in W, a, b \in R$

It is clear that:  $(W^2, +, \cdot)$  is a commutative ring, and  $(W^2, +, \times)$  is a module over W, it is called the weak fuzzy complex Euclidean module with two dimensions.

### Definition:

Let  $W^3 = \{(x, y, z); x, y, z \in W\}$ , define:

$$(+)\ W^3 \times W^3 \rightarrow W^3; (x_0, x_1, x_2) + (y_0, y_1, y_2) = (x_0 + y_0, x_1 + y_1, x_2 + y_2)$$

$$(\cdot)\ W^3 \times W^3 \rightarrow W^3; (x_0, x_1, x_2) \cdot (y_0, y_1, y_2) = (x_0 y_0, x_1 y_1, x_2 y_2)$$

$$(\times)\ W \times W^3 \rightarrow W^3; (a + bJ) \times (x_0, x_1, x_2) = ((a + bJ)x_0, (a + bJ)x_1, (a + bJ)x_2)$$

It is clear that:  $(W^3, +, \cdot)$  is a commutative ring, and  $(W^3, +, \times)$  is a module over W, it is called the weak fuzzy complex Euclid module with three dimensions.

### Remark:

Consider  $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}, \mathbb{R}^2 \times \mathbb{R}^2, \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$ , and the following algebraic operations:

$$(+)\ \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2; (x_0, x_1) + (y_0, y_1) = (x_0 + y_0, x_1 + y_1)$$

$$(\cdot)\ \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2; (x_0, x_1) \cdot (y_0, y_1) = (x_0 y_0, x_1 y_1)$$

$$(\times)\ \mathbb{R}^2 \times (\mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2; (a, b) \times ((x, y), (z, t)) = ((ax, by), (az, bt)),$$

$$(+)\ (\mathbb{R}^2 \times \mathbb{R}^2) \times (\mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2;$$

$$((x_0, x_1), (y_0, y_1)) + ((x'_0, x'_1), (y'_0, y'_1)) = ((x_0 + x'_0, x_1 + x'_1), (y_0 + y'_0, y_1 + y'_1))$$

$$(\cdot)\ (\mathbb{R}^2 \times \mathbb{R}^2) \times (\mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2;$$

$$((x_0, x_1), (y_0, y_1)) \cdot ((x'_0, x'_1), (y'_0, y'_1)) = ((x_0 x'_0, x_1 x'_1), (y_0 y'_0, y_1 y'_1))$$

It is known that  $(\mathbb{R}^2, +, \cdot)$  is a commutative ring,  $(\mathbb{R}^2 \times \mathbb{R}^2, +, \cdot)$  is a commutative ring,  $(\mathbb{R}^2 \times \mathbb{R}^2, +, \times)$  is a module over  $\mathbb{R}^2$ .

Also,  $(+)\ (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2) \times (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2;$

$$((x_0, x_1), (y_0, y_1), (z_0, z_1)) + ((x'_0, x'_1), (y'_0, y'_1), (z'_0, z'_1)) = ((x_0 + x'_0, x_1 + x'_1), (y_0 + y'_0, y_1 + y'_1), (z_0 + z'_0, z_1 + z'_1))$$

$$(\cdot)\ (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2) \times (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2;$$

$$((x_0, x_1), (y_0, y_1), (z_0, z_1)) \cdot ((x'_0, x'_1), (y'_0, y'_1), (z'_0, z'_1)) = ((x_0 x'_0, x_1 x'_1), (y_0 y'_0, y_1 y'_1), (z_0 z'_0, z_1 z'_1))$$

$$(\times)\ \mathbb{R}^2 \times (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2) \rightarrow \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2;$$

$$(a, b) \times ((x_0, x_1), (y_0, y_1), (z_0, z_1)) = ((ax_0, bx_1), (ay_0, by_1), (az_0, bz_1))$$

It is clear that  $(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2, +, \cdot)$  is a commutative ring,  $(\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2, +, \times)$  is a module over  $\mathbb{R}^2$ .

**Theorem:**

The ring  $(W, +, \cdot)$  is isomorphic to  $(\mathbb{R}^2, +, \cdot)$ .

Proof:

Define:  $T_1: W \rightarrow \mathbb{R}^2$  such that:

$$T_1(x_0, x_1J) = (x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t}); J^2 = t \in ]0, 1[$$

It is clear that  $T_1$  is well defined mapping, and for  $X = x_0 + x_1J, Y = y_0 + y_1J \in W$ , we have:

$$X + Y = (x_0 + y_0) + (x_1 + y_1)J, X \cdot Y = (x_0y_0 + x_1y_1t) + J(x_0y_1 + x_1y_0),$$

$$T_1(X + Y) = (C_0, C_1); C_0 = (x_0 + y_0) - \sqrt{t}(x_1 + y_1) = (x_0 - \sqrt{t}x_1) + ((y_0 - \sqrt{t}y_1),$$

$$C_1 = (x_0 + y_0) + \sqrt{t}(x_1 + y_1) = (x_0 + \sqrt{t}x_1) + ((y_0 + \sqrt{t}y_1), \text{ thus}$$

$$T_1(X + Y) = T_1(X) + T_1(Y).$$

$$T_1(XY) = (D_0, D_1); D_0 = x_0y_0 + x_1y_1t - \sqrt{t}(x_0y_1 + x_1y_0) = (x_0 - \sqrt{t}x_1)(y_0 - \sqrt{t}y_1), D_1 = x_0y_0 + x_1y_1t + \sqrt{t}(x_0y_1 + x_1y_0) = (x_0 + \sqrt{t}x_1)(y_0 + \sqrt{t}y_1), \text{ thus } T_1(XY) = T_1(X)T_1(Y).$$

$$\text{Assume that } T_1(X) = 0, \text{ then } \begin{cases} x_0 - x_1\sqrt{t} = 0 \\ x_0 + x_1\sqrt{t} = 0 \end{cases} \text{ so } x_0 = x_1 = 0$$

And  $\ker(T_1) = \{0\}$ , which means that  $T_1$  is injective.

$\text{Im}(T_1) = \mathbb{R}^2$ , hence  $T_1$  is surjective and then it is a bijection.

This implies the proof.

Remark:

$$T_1^{-1}\mathbb{R}^2 \rightarrow W; T_1^{-1}(x, y) = \frac{1}{2}(x + y) + \frac{1}{2\sqrt{t}}J(y - x); J^2 = t \in ]0, 1[.$$

**Definition:**

Let  $X = x_0 + x_1J, Y = y_0 + y_1J \in W$ , we say that  $X \geq Y$  if and only if

$$\begin{cases} x_0 - \sqrt{t}x_1 \geq y_0 - \sqrt{t}y_1 \\ x_0 + \sqrt{t}x_1 \geq y_0 + \sqrt{t}y_1 \end{cases}$$

**Example:**

Consider  $X = 4 + J, Y = 1 + 2J$  with  $J^2 = t = \frac{1}{4} \in ]0, 1[$ :

$$x_0 - \sqrt{t}x_1 = \frac{7}{2} \geq y_0 - \sqrt{t}y_1 = 0$$

$$x_0 + \sqrt{t}x_1 = \frac{9}{2} \geq y_0 + \sqrt{t}y_1 = 2, \text{ thus } X \geq Y.$$

**Definition.**

1]  $X = x_0 + x_1J \in W$  is called positive weak fuzzy complex number if  $X \geq 0$ , i.e:  $x_0 - \sqrt{t}x_1 \geq 0, x_0 + \sqrt{t}x_1 \geq 0$ .

2]  $X$  is negative if  $X \leq 0$ , i.e:  $x_0 - \sqrt{t}x_1 \leq 0, x_0 + \sqrt{t}x_1 \leq 0$ .

**Theorem:**

$(\geq)$  is a partial order relation.

Proof:

Let  $X = x_0 + x_1J, Y = y_0 + y_1J, Z = z_0 + z_1J \in W$ ,

$$X \geq X \text{ that is because: } \begin{cases} x_0 - x_1\sqrt{t} \geq x_0 - \sqrt{t}x_1 \\ x_0 + x_1\sqrt{t} \geq x_0 + \sqrt{t}x_1 \end{cases}$$

$$\text{If } X \geq Y \text{ and } Y \geq X, \text{ then: } \begin{cases} x_0 - x_1\sqrt{t} \geq y_0 - \sqrt{t}y_1 \\ x_0 + x_1\sqrt{t} \geq y_0 + \sqrt{t}y_1 \end{cases} \text{ and}$$

$$\begin{cases} y_0 - \sqrt{t}y_1 \geq x_0 - \sqrt{t}x_1 \\ y_0 + \sqrt{t}y_1 \geq x_0 + \sqrt{t}x_1 \end{cases}, \text{ this implies that:}$$

$$\begin{cases} x_0 - x_1\sqrt{t} \geq y_0 - \sqrt{t}y_1 \\ x_0 + x_1\sqrt{t} \geq y_0 + \sqrt{t}y_1 \end{cases}, \text{ thus } x_0 = y_0, x_1 = y_1 \text{ and } X = Y,$$

If  $X \geq Y$  and  $Y \geq Z$ , then:

$$\begin{cases} x_0 - x_1\sqrt{t} \geq y_0 - y_1\sqrt{t} \geq z_0 - z_1\sqrt{t} \\ x_0 + x_1\sqrt{t} \geq y_0 + y_1\sqrt{t} \geq z_0 + z_1\sqrt{t} \end{cases}$$

Thus  $X \geq Z$ , and the proof is complete.

**Definition:**

We define the absolute value of  $X = x_0 + x_1J \in W$  as follows,

$$|X| = \frac{1}{2} [|x_0 + \sqrt{t}x_1| + |x_0 - \sqrt{t}x_1|] + \frac{1}{2\sqrt{t}} J[|x_0 + \sqrt{t}x_1| - |x_0 - \sqrt{t}x_1|]$$

**Example:**

For  $J^2 = t = \frac{1}{9}$ , consider:  $X = -3 + 6J$ , we have:

$$\begin{cases} x_0 - \sqrt{t}x_1 = -3 - \frac{6}{3} = -5 \\ x_0 + \sqrt{t}x_1 = -3 + \frac{6}{3} = -1 \end{cases}$$

$$\Rightarrow |X| = \frac{1}{2}(|-5| + |-1|) + \frac{1}{(2)(\frac{1}{9})}J[|-1| - |-5|] = 3 + \frac{3}{2}J(-4) = 3 - 6J.$$

**Remark:**

$|X| \geq 0$  according to the partial order relation defined on  $W$  for all  $X \in W$

**Theorem:**

$$1) (W^2, +, \cdot) \cong (\mathbb{R}^2 \times \mathbb{R}^2, +, \cdot)$$

$$2) (W^3, +, \cdot) \cong (\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2, +, \cdot)$$

Proof:

1) Define  $T_2: W^2 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2$  such that:

$$T_2(X, Y) = (T_1(x), T_1(y)); x, y \in W$$

$T_2$  is well defined, that is because:

If  $(x_0, y_0) = (x_1, y_1); x_0, x_1, y_0, y_1 \in W$ , then:

$$\begin{cases} x_0 = x_1 \\ y_0 = y_1 \end{cases} \text{ so that } (T_1(x_0), T_1(y_0)) = (T_1(x_1), T_1(y_1))$$

$$\text{Hence, } T_2(x_0, y_0) = T_2(x_1, y_1).$$

$$\ker(T_2) = \{(x, y) \in W^2; T_2(x, y) = ((0, 0), (0, 0))\}, \text{ thus}$$

$$\begin{cases} T_1(x) = (0, 0) \\ T_1(y) = (0, 0) \end{cases} \text{ which implies that } X=Y=0 \text{ and } \ker(T_2) = \{(0, 0)\}.$$

Also,  $\text{Im}(T_2) = (\text{Im}(T_1), \text{Im}(T_1)) = \mathbb{R}^2 \times \mathbb{R}^2$ , thus  $T_2$  is a bijection for  $(x_0, y_0), (x_1, y_1) \in W^2$ , we have:

$$\begin{aligned} T_2[(x_0, y_0) + (x_1, y_1)] &= T_2[(x_0 + x_1, y_0 + y_1)] \\ &= (T_1(x_0 + x_1), T_1(y_0 + y_1)) = (T_1(x_0), T_1(y_0)) + (T_1(x_1), T_1(y_1)) \\ &= T_2(x_0, y_0) + T_2(x_1, y_1). \end{aligned}$$

$$\begin{aligned} T_2[(x_0, y_0) \cdot (x_1, y_1)] &= T_2[(x_0x_1, y_0y_1)] = (T_1(x_0x_1), T_1(y_0y_1)) = (T_1(x_0), T_1(y_0)) \cdot (T_1(x_1), T_1(y_1)) = \\ &= T_2(x_0, y_0) \cdot T_2(x_1, y_1) \end{aligned}$$

Thus  $T_2$  is a ring isomorphism.

2) Define  $T_3: W^3 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$  such that:

$$T_3(X, Y, Z) = (T_1(x), T_1(y), T_1(z)); x, y, z \in W$$

$T_3$  can be proved that it is a ring isomorphism by a similar argument of  $T_2$ .

**Example:**

Let  $A = (3 + 2J, 1 - 4J) \in W^2; J^2 = t = \frac{1}{4} \in ]0, 1[$ , then

$$T_2(A) = (T_1(3 + 2J), T_1(1 - 4J)) = ((2, 4), (3, -1)).$$

$$\text{Remark that } T_1(3 + 2J) = \left(3 - \left(\frac{1}{2}\right)(2), 3 + \left(\frac{1}{2}\right)(2)\right) = (2, 4)$$

$$T_1(1 - 4J) = \left(1 + \frac{1}{2}(4), 1 + \frac{1}{2}(-4)\right) = (3, -1)$$

Let  $B = (3 + 2J, 1 - 4J, 3 + 2J) \in W^3$ , then:

$$T_3(B) = (T_1(3 + 2J), T_1(1 - 4J), T_1(3 + 2J)) = ((2, 4), (3, -1), (2, 4))$$

**Remark:**

1) the inverse isomorphism of  $T_2$  is:  $T_2^{-1}: \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow W^2$  such that:

$$\begin{aligned} T_2^{-1}((x_0, x_1), (y_0, y_1)) &= (T_1^{-1}(x_0, x_1), T_1^{-1}(y_0, y_1)) \\ &= \left(\frac{x_0 + x_1}{2} + \frac{J}{2\sqrt{t}}(x_1 - x_0), \frac{y_0 + y_1}{2} + \frac{J}{2\sqrt{t}}(y_1 - y_0)\right) \end{aligned}$$

2) the inverse isomorphism of  $T_3$  is:  $T_3^{-1}: \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow W^3$  such that:

$$T_3^{-1}((x_0, x_1), (y_0, y_1), (z_0, z_1)) = (T_1^{-1}(x_0, x_1), T_1^{-1}(y_0, y_1), T_1^{-1}(z_0, z_1))$$

**Definition: [15-16]**

Let  $M$  be a module over the ring  $R$ ,  $N$  be a module over  $T$ , then  $f: M \rightarrow N$  such that:  $f(x + y) = f(x) + f(y)$ ,  $f(a \cdot x) = g(a)f(x)$ ;  $g: R \rightarrow T$  is called a semi module isomorphism if and only if:

1)  $f$  is a bijection.

2)  $g$  is a ring isomorphism.

Theorem:

1) the module  $(W^2, +, \times)$  is semi isomorphic to  $(R^2 \times R^2, +, \times)$ .

2) the module  $(W^3, +, \times)$  is semi isomorphic to  $(R^2 \times R^2 \times R^2, +, \times)$ .

Proof:

1] Consider  $T_2: W^2 \rightarrow R^2 \times R^2$ ,  $T_1: W \rightarrow R^2$ , we have  $T_2$  is a bijection that preserves addition, i.e:

$$T_2[(x_0, x_1) + (y_0, y_1)] = T_2(x_0, x_1) + T_2(y_0, y_1)$$

Also,  $T_1$  is a ring isomorphism between  $W$  and  $R^2$ .

Now, we must prove that  $T_2[A \cdot (x, y)] = T_1(A) \cdot T_2(x, y)$  for all  $(x, y) \in W^2$ ,  $A \in W$

$$\begin{aligned} T_2[A \cdot (x, y)] &= T_2(Ax, Ay) = (T_1(Ax), T_1(Ay)) \\ &= (T_1(A)T_1(x), T_1(A)T_1(y)) = T_1(A) \cdot (T_1(x), T_1(y)) = T_1(A) \cdot T_2(x, y) \end{aligned}$$

This means that  $T_2$  is a semi module isomorphism.

2] It can be proved by a similar argument by considering

$$T_3: W^3 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \text{ and } T_1: W \rightarrow \mathbb{R}^2$$

The classification of A-curves  $C_{A_t}$ ;  $t \in ]0, 1[$

**Definition: [7]**

1] The  $A_t$ -Curve in  $\mathbb{R}^2$  is defined by the following Vectorial equation:

$$\begin{cases} \|x\|^2 + t \cdot \|y\|^2 = r_1; \\ x_1x_2 + y_1y_2 = r_2 \end{cases}; X = (x_1, y_1), Y = (x_2, y_2) \in \mathbb{R}^2, t \in ]0, 1[$$

$$r_1, r_2 \geq 0$$

2] The  $A_t$ -Curve in  $\mathbb{R}^3$  is defined by the following Vectorial equation:

$$\begin{cases} \|x\|^2 + t \cdot \|y\|^2 = r_1 \\ x_1x_2 + y_1y_2 + z_1z_2 = r_2 \end{cases}; X = (x_1, y_1, z_1), Y = (x_2, y_2, z_2) \in \mathbb{R}^3,$$

$$t \in ]0, 1[, r_1, r_2 \geq 0$$

**Theorem: [7]**

1]  $C_{A_t}$  in  $\mathbb{R}^2$  is equivalent to a weak fuzzy complex circle:  $X^2 + Y^2 = r_1 + 2r_2J$ ;  $J^2 = t$ ,  $X = x_1 + x_2J$ ,  $Y = y_1 + y_2J \in W$

2]  $C_{A_t}$  in  $\mathbb{R}^3$  is equivalent to a weak fuzzy complex sphere:  $X^2 + Y^2 + Z^2 = r_1 + 2r_2J$ ;

$$X = x_1 + x_2J, Y = y_1 + y_2J, Z = z_1 + z_2J \in W$$

**Theorem:**

1]  $C_{A_t}$  in  $\mathbb{R}^2$  is isomorphic to the cartesian product to two circles in the Euclidean space  $\mathbb{R}^2$ , i.e: for every single point  $(x', y') \in C_{A_t}$  there exists only two points (up to isomorphism) and each point belongs to an ordinary Euclidean circle, if  $r_1 - 2r_2\sqrt{t} > 0$ .

Also, it is isomorphic to an empty set if  $r_1 - 2r_2\sqrt{t} < 0$ , and it is isomorphic to a single point if  $r_1 = r_2 = 0$ .

1]  $C_{A_t}$  in  $\mathbb{R}^3$  is isomorphic to the cartesian product of two classical Euclidean spheres if  $r_1 - 2r_2\sqrt{t} > 0$ , and to an empty set if  $r_1 - 2r_2\sqrt{t} < 0$  and to a single point if  $r_1 = r_2 = 0$ .

Proof:

1] the equation of  $C_{A_t}$  in  $\mathbb{R}^2$  is:

$$X^2 + Y^2 = r_1 + 2r_2J; X = x_1 + x_2J, Y = y_1 + y_2J, J^2 = t \in ]0, 1[.$$

By applying the isomorphism  $T_1$ , we get:

$$T_1(X^2) + T_1(Y^2) = T_1(r_1 + 2r_2J), \text{ hence:}$$

$$[T_1(x)]^2 + [T_1(y)]^2 = T_1(r_1 + 2r_2J), \text{ thus:}$$

$$\begin{cases} (x_1 - x_2\sqrt{t})^2 + (y_1 - y_2\sqrt{t})^2 = r_1 - 2r_2J \\ (x_1 + x_2\sqrt{t})^2 + (y_1 + y_2\sqrt{t})^2 = r_1 + 2r_2J \end{cases}$$

We change the variables as follows:

$$\begin{cases} x_1 - x_2\sqrt{t} = X_1 \\ x_1 + x_2\sqrt{t} = X_2 \end{cases} \begin{cases} y_1 - y_2\sqrt{t} = Y_1 \\ y_1 + y_2\sqrt{t} = Y_2 \end{cases}$$

Then,

$$\begin{cases} X_1^2 + Y_1^2 = r_1 - 2r_2\sqrt{t} \\ X_2^2 + Y_2^2 = r_1 + 2r_2\sqrt{t} \end{cases}$$

According to the assumption, we have  $r_1, r_2 \geq 0$ , this implies the following cases,

Case 1:

If  $r_1 = r_2 = 0$ , then  $C_{A_t}$  has only one point (0,0).

Case 2:

If  $r_1 - 2r_2\sqrt{t} < 0$ , then  $C_{A_t}$  is an empty set

Case 3:

If  $r_1 - 2r_2\sqrt{t} > 0$ , then  $C_{A_t}$  is isomorphic to the direct product of the two classical circles:

$$\begin{cases} X_1^2 + Y_1^2 = r_1 - 2r_2\sqrt{t} \\ X_2^2 + Y_2^2 = r_1 + 2r_2\sqrt{t} \end{cases}$$

2]  $C_{A_t}$  in  $\mathbb{R}^3$  has the following equation:

$$X^2 + Y^2 + Z^2 = r_1 + 2r_2J; J^2 = t \in ]0,1[ \quad X = x_1 + x_2J, \quad Y = y_1 + y_2J, \quad Z = z_1 + z_2J \in W.$$

By using  $T_1$ , we get:

$$T_1(X^2) + T_1(Y^2) + T_1(Z^2) = T_1(r_1 + 2r_2J), \text{ thus:}$$

$$\begin{cases} (x_1 - \sqrt{t}x_2)^2 + (y_1 - \sqrt{t}y_2)^2 + (z_1 - \sqrt{t}z_2)^2 = r_1 - 2r_2\sqrt{t} \\ (x_1 + \sqrt{t}x_2)^2 + (y_1 + \sqrt{t}y_2)^2 + (z_1 + \sqrt{t}z_2)^2 = r_1 + 2r_2\sqrt{t} \end{cases}$$

If  $r_1 - 2r_2\sqrt{t} > 0$ , we get two classical Euclidean spheres.

If  $r_1 = r_2 = 0$ , we get a single point  $(0,0)$ .

If  $r_1 - 2r_2\sqrt{t} < 0$ , we get an empty set

#### Example:

Consider the following  $C_{A_t}$ -Curve in  $\mathbb{R}^2$

$$\begin{cases} \|x\|^2 + \|y\|^2 t = 4 \\ x_1x_2 + y_1y_2 = 2 \end{cases} \text{ with } t = \frac{1}{4} \in ]0,1[; \quad X = (x_1, y_1), Y = (x_2, y_2) \in \mathbb{R}^3,$$

With the following representation:

$$X_J^2 + Y_J^2 = 4 + 4J(C_J); \quad X_J = x_1 + x_2J, \quad Y_J = y_1 + y_2J$$

We have  $r_1 = 4, r_2 = 2, r_1 + 2r_2\sqrt{t} = 6, r_1 - 2r_2\sqrt{t} = 2 > 0$ , Hence  $C_{A_{\frac{1}{4}}}$  is equivalent to the following classical circles:

$$\begin{cases} X_1^2 + Y_1^2 = 2(C_1) \\ X_2^2 + Y_2^2 = 6(C_2) \end{cases}; \quad X_1, Y_1, X_2, Y_2 \in \mathbb{R}, \text{ and } X_1 = x_1 - x_2\sqrt{t}$$

$$Y_2 = y_1 + y_2\sqrt{t}, \quad X_2 = x_1 + x_2\sqrt{t}, \quad Y_1 = y_1 - y_2\sqrt{t}$$

For example:  $(1,1) \in C_1, (\sqrt{6}, 0) \in C_2$ , i.e:

$$\begin{aligned} x_1 = 1, x_2 = \sqrt{6}, y_1 = 1, y_2 = 0, \text{ this generates a point in } \mathbb{R}^2 \times \mathbb{R}^2 \text{ as follows:} \\ = ((x_1, x_2), (y_1, y_2)) = ((1, \sqrt{6}), (1, 0)) \end{aligned}$$

The inverse image  $T_2^{-1}(L)$  general a point on  $C_J$  as follows

$$T_2^{-1}(L) = \left( \frac{1 + \sqrt{6}}{2} + J(\sqrt{6} - 1), \frac{1}{2} - J \right) \in C_J$$

$$\text{Remark that: } \left( \frac{1 + \sqrt{6}}{2} + J(\sqrt{6} - 1) \right)^2 + \left( \frac{1}{2} - J \right)^2 = 4 + 4J$$

This point gives us a solution to the original vectorial equation  $\begin{cases} \|x\|^2 + \|y\|^2 t = 4 \\ x_1x_2 + y_1y_2 = 2 \end{cases}; \quad t = \frac{1}{4}$

$$\text{This solution } X = (x_1, y_1) = \left( \frac{1 + \sqrt{6}}{2}, \frac{1}{2} \right), \quad Y = (x_2, y_2) = (\sqrt{6} - 1, -1)$$

#### Example:

Let's find a solution of the vectorial equation:

$$\begin{cases} \|x\|^2 + t \cdot \|y\|^2 = 9 \\ x_1x_2 + y_1y_2 + z_1z_2 = 6 \end{cases}; \quad X = (x_1, y_1, z_1), Y = (x_2, y_2, z_2) \in \mathbb{R}^3 \quad t = \frac{1}{9} \in ]0,1[.$$

Any solution to the previous vectorial equation is equivalent to a single point on  $C_{A_{\frac{1}{9}}}$  in  $\mathbb{R}^3$ .

The weak fuzzy complex formula of it is:

$$C_J)X_J^2 + Y_J^2 + Z_J^2 = 9 + 12J; \quad X_J^2 = x_1 + x_2J, \quad Y_J^2 = y_1 + y_2J, \quad Z_J^2 = z_1 + z_2J \in W \text{ on the other hand, we have}$$

$$r_1 = 9, r_2 = 6, r_1 + 2r_2\sqrt{t} = 9 + 2(6)\left(\frac{1}{3}\right) = 13, r_1 - 2r_2\sqrt{t} = 9 - 2(6)\left(\frac{1}{3}\right) = 5 > 0, \text{ hence } C_J \text{ can be}$$

represented by the following classical spheres:

$$\begin{cases} X_1^2 + Y_1^2 + Z_1^2 = 5(C_1) \\ X_2^2 + Y_2^2 + Z_2^2 = 13(C_2) \end{cases}, \text{ where } \begin{cases} X_1 = x_1 - x_2\sqrt{t} = x_1 - \frac{x_2}{3} \in \mathbb{R} \\ Y_1 = y_1 - y_2\sqrt{t} = y_1 - \frac{y_2}{3} \in \mathbb{R} \\ Z_1 = z_1 - z_2\sqrt{t} = z_1 - \frac{z_2}{3} \in \mathbb{R} \end{cases}$$

$$\text{And: } \begin{cases} X_2 = x_1 + x_2\sqrt{t} = x_1 + \frac{x_2}{3} \in \mathbb{R} \\ Y_2 = y_1 + y_2\sqrt{t} = y_1 + \frac{y_2}{3} \in \mathbb{R} \\ Z_2 = z_1 + z_2\sqrt{t} = z_1 + \frac{z_2}{3} \in \mathbb{R} \end{cases}$$

$(2,1,0) \in C_1, (3,0,2) \in C_2$ , hence:

$$L = ((x_1, x_2), (y_1, y_2), (z_1, z_2)) = ((2,3), (1,0), (0,2)) \in R^2 \times R^2 \times R^2$$

The inverse image  $T_3^{-1}(L) = \left( \frac{5}{2} + \frac{3}{2}J, \frac{1}{2} - \frac{3}{2}J, 1 + 3J \right)$  is a point on  $(C_J)$ .

Thus:  $X = (x_1, y_1, z_1) = \left( \frac{5}{2}, \frac{1}{2}, 1 \right), Y = (x_2, y_2, z_2) = \left( \frac{3}{2}, -\frac{3}{2}, 3 \right)$  is a solution to the vectorial equation.

### 3. Conclusion

In this article, we have studied the classification problem of A-curves generated from the Euclidean vectorial equation  $\begin{cases} \|x\|^2 + t\|y\|^2 = r \\ x_1x_2 + y_1y_2 + z_1z_2 = k \end{cases}$ ; by using some novel suggested semi-module homomorphisms and semi-module isomorphisms that helped in formulating weak fuzzy complex numbers as real vectors. Also, we have presented many examples to explain the contribution of our work.

In the future, we aim to classify weak fuzzy complex inner product spaces by using the same semi-homomorphisms.

### References

- [1] Deckelman, S., Robson, B. Split-Complex Numbers and Dirac Bra-kets., *Communications In Information and Systems*, **14**, (2014), 135-159.
- [2] Akar, M., Yuce, S., Sahin, S., On the Dual Hyperbolic Numbers and The Complex Hyperbolic Numbers, *JCSCM*, **8**, (2018), DOI: 10.20967/jcscm.2018.01.001.
- [3] Hatip, A., An Introduction To Weak Fuzzy Complex Numbers, *Galoitica Journal Of Mathematical Structures and Applications*, **3**, (2023).
- [4] Ali, R., On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces, *Neoma Journal Of Mathematics and Computer Science*, (2023).
- [5] Hatip, A., On The Weak Fuzzy Complex Vector Spaces, *Galoitica Journal Of Mathematical Structures And Applications*, **3**, (2023).
- [6] Ali, R., A Short Note On The Solution of n-Refined Neutrosophic Linear Diophantine Equations, *International Journal Of Neutrosophic Science*, **15**, (2021).
- [7] Alhasan, Y., Alfahal, A., Abdulfatah, R., Nordo, G. & Zahra, M., On Some Novel Results About Weak Fuzzy Complex Matrices. *International Journal of Neutrosophic Science*, **21**, (2023), 134-140. <https://doi.org/10.54216/IJNS.210112>
- [8] Ahmad, K., Bal, M., and Aswad, M., A Short Note On The Solutions Of Fermat's Diophantine Equation In Some Neutrosophic Rings, *Journal Of Neutrosophic and Fuzzy Systems*, (2022).
- [9] Abobala, M., Partial Foundation of Neutrosophic Number Theory, *Neutrosophic Sets and Systems*, **39**, (2021).
- [10] Alfahal, A., Abobala, M., Alhasan, Y., and Abdulfatah, R., Generating Weak Fuzzy Complex and Anti Weak Fuzzy Complex Integer Solutions for Pythagoras Diophantine Equation  $X^2 + Y^2 = Z^2$ , *International Journal of Neutrosophic Science*, **22**, (2023).
- [11] Alhasan, Y., Xu, L., Abdulfatah, R., & Alfahal, A., The Geometrical Characterization for The Solutions of a Vectorial Equation By Using Weak Fuzzy Complex Numbers and Other Generalizations Of Real Numbers. *International Journal of Neutrosophic Science (IJNS)*, **21**, (2023), 155-159. <https://doi.org/10.54216/IJNS.210415>
- [12] Galarza, F., Flores, M., Rivero, D., & Abobala, M., On Weak Fuzzy Complex Pythagoras Quadruples. *International Journal of Neutrosophic Science (IJNS)*, **22**, (2023), 108-113. <https://doi.org/10.54216/IJNS.220209>.
- [13] Razouk, L., Mahmoud, S., & Ali, M., A Computer Program For The System Of Weak Fuzzy Complex Numbers And Their Arithmetic Operations Using Python. *Galoitica: Journal Of Mathematical Structures and Applications*, **8**, (2023), 45-51. <https://doi.org/10.54216/GJMSA.080104>.
- [14] Merkepci, M., and Abobala, M., " On Some Novel Results About Split-Complex Numbers, The Diagonalization Problem And Applications To Public Key Asymmetric Cryptography", *Journal of Mathematics*, Hindawi, 2023.
- [15] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", *International journal of neutrosophic science*, 2022.
- [16] Abobala, M., "Semi Homomorphisms and Algebraic Relations Between Strong Refined Neutrosophic Modules and Strong Neutrosophic Modules", *Neutrosophic Sets and Systems*, 2021.
- [17] Murtada Ali Maqdisi. (2023). The Decay of The Solutions of a Nonlinear Viscoelastic Hyperbolic Equation. *Pure Mathematics for Theoretical Computer Science*, **1** ( 1 ), 47-55 (Doi : <https://doi.org/10.54216/PMTCS.010104>)